



Brain and Other Central Nervous System (CNS) Cancer


A Fact Sheet from the North Carolina Central Cancer Registry, State Center for Health Statistics

August 2017


Cancer of the brain and other central nervous system (CNS), excluding benign cases, was the 17th most frequently occurring and the ninth leading cause of cancer death in North Carolina from 2010 to 2014. It is anticipated that 774 people (408 males and 336 females) in North Carolina will be diagnosed with and 534 people (297 males and 237 females) will die of cancer of the brain and other CNS in 2017.

Incidence


The percentage of cases of brain and other CNS cancer from 2010 to 2014 is displayed by age group in Figure 1. About 12 percent of brain and other CNS cancer were diagnosed in people ages 0 to 19.



Between 2010 and 2014, the age-adjusted incidence rate for brain and other CNS cancer in North Carolina was 6.5 per 100,000 persons per year. In general, men are more likely to be diagnosed with brain and other CNS cancer than women (Figure 2).




From 2003 to 2014, brain and other CNS cancer incidence rates have remained fairly stable for both men and women (Figure 3).




Stage at Diagnosis*

Figure 4 shows the stage distribution of brain and other CNS cancer cases diagnosed between 2010 and 2014. More than 75 percent of brain and other CNS cancer cases were diagnosed at the localized stage.



Mortality

Between 2010 and 2014, the percentage of brain and other CNS cancer deaths is displayed by age group in Figure 5. People ages 65 to 84 had the highest percentage of deaths compared with other age groups.




Data Sources and Methods

Data on North Carolina cases were obtained from the North Carolina Central Cancer Registry (CCR). Hospitals are the primary source of data. The CCR supplements hospital data with reports from physicians who diagnose cases in a non-hospital setting. The CCR also collects data from pathology laboratories and freestanding treatment centers. Data on cancer deaths were obtained from Statistical Services in the State Center for Health Statistics. Population data from the National Center for Health Statistics were used in the denominators of the rates, which are expressed per 100,000 persons. Rates were age-adjusted using the 2000 United States Census data. To examine trends, three-year overlapping rates were used to improve stability over time. Stage at diagnosis was defined according to Surveillance, Epidemiology, and End Results Summary Stage guidelines as in situ, localized, regional, distant and unknown/NA. For further information about the North Carolina CCR, please visit www.schs.state.nc.us/units/ccr.

* According to the National Cancer Institute (NCI), "many cancer registries, such as NCI's Surveillance, Epidemiology, and End Results Program (SEER), use summary staging. This system is used for all types of cancer. It groups cancer cases into five main categories: **In situ**—Abnormal cells are present only in the layer of cells in which they developed. **Localized**—Cancer is limited to the organ in which it began, without evidence of spread. **Regional**—Cancer has spread beyond the primary site to nearby lymph nodes or organs and tissues. **Distant**—Cancer has spread from the primary site to distant organs or distant lymph nodes. **Unknown**—There is not enough information to determine the stage." Additional information on staging can be found at www.cancer.gov/cancertopics/factsheet/detection/staging.

The age-adjusted mortality rate of brain and other CNS cancer from 2010 to 2014 was 4.5 per 100,000 persons per year. In general, men are more likely to die from brain and other CNS cancer than women (Figure 6).



From 2003 to 2014, brain and other CNS cancer mortality rates remained fairly stable for both men and women (Figure 7).

